

Banixin-50

Flunexine 5% injection



*For treatment of acute inflammation reactions
associated with infectious diseases.*


Banixin-50

Solution for parenteral administration

Flunixin is a non-steroidal anti-inflammatory drug (NSAID), and a non-narcotic analgesic with antipyretic properties. It is used in the alleviation of inflammation and pain associated with musculoskeletal disorders and colic. In addition it is used for the control of acute inflammation reactions associated with infectious diseases. Flunixin is also applied as adjunctive therapy in the treatment of respiratory diseases.

CONTAINS PER ML

Flunixin (as meglumine) 50 mg.



INDICATIONS

Cattle:

Banixin-50 is indicated for the control of pyrexia associated with bovine respiratory disease, endotoxemia and acute bovine mastitis. Banixin-50 is also indicated for the control of inflammation in endotoxemia and as an adjunct to antimicrobial therapy to reduce clinical signs of acute inflammation in cases of infectious respiratory disease.

Horses:

Banixin-50 is indicated for the alleviation of inflammatory signs associated with musculoskeletal disorders and for the alleviation of visceral pain associated with colic.

Swine:

Banixin-50 is indicated as an adjunctive therapy in the treatment of swine respiratory diseases.

DOSAGE

For intramuscular or intravenous administration.

Cattle:

2 ml per 45 kg body weight. Repeat if necessary at 24 h intervals for up to 3 consecutive days.

Swine:

2 ml per 45 kg body weight once by intramuscular injection only.

Horses:

1 ml per 45 kg body weight by intravenous injection only. Treatment may be repeated once or twice if colic recurs. In case of musculo-skeletal disorders, treatment is once daily for up to 5 days according to clinical response.

Inject the product slowly, and apply at body temperature. Administration should be stopped immediately if signs of intolerance occur. An adequate water supply should be provided.

WITHDRAWAL TIMES

- For meat:

Cattle : 14 days.
Pigs : 24 days.

- For milk:

Cattle : 2 days.

Do not use in horses producing meat or milk for human consumption.

PACKAGING

Vial of 50 ml.



sales@interchemie.com | interchemie.com



Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XFIFLS-EN-01-10092020