

Interflox-100

Enrofloxacin 10% injection



*For the treatment of Gastrointestinal and
respiratory infection caused by enrofloxacin
sensitive micro-organisms*

Interflox-100

Solution for parenteral administration

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like *Campylobacter*, *E. coli*, *Haemophilus*, *Pasteurella*, *Mycoplasma* and *Salmonella* spp.

CONTAINS PER ML

Enrofloxacin 100 mg.

INDICATIONS

Gastrointestinal and respiratory infections caused by enrofloxacin sensitive micro-organisms, like *Campylobacter*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella* and *Salmonella* spp. in calves, cattle, sheep, goats and swine.

DOSAGE

For intramuscular or subcutaneous administration.

Calves, cattle, sheep and goats:

1 ml per 20 - 40 kg body weight for 3 - 5 days.

Swine:

1 ml per 20 - 40 kg body weight for 3 - 5 days.

WITHDRAWAL TIMES

- For meat:
 - Calves, cattle, sheep and goats : 21 days.
 - Swine : 14 days.
- For milk : 4 days.

PACKAGING

Vial of 50 and 100 ml.



sales@interchemie.com | interchemie.com



Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XF1EN10-EN-01-27082020