

Interspectin-L

Spectinomycin & Lincomycin injection



*Treatment of Gastrointestinal and
respiratory infections caused by lincomycin
and spectinomycin sensitive micro-organisms*

Interspectin-L

Solution for parenteral administration

The combination of lincomycin and spectinomycin acts additive and in some cases synergistic. Spectinomycin acts bacteriostatic or bactericidal, depending on the dose, against mainly Gram-negative bacteria like *Campylobacter*, *E. coli* and *Salmonella* spp. Lincomycin acts bacteriostatic against mainly Gram-positive bacteria like *Mycoplasma*, *Treponema*, *Staphylococcus* and *Streptococcus* spp. Cross-resistance of lincomycin with macrolides can occur.

CONTAINS PER ML

Spectinomycin base 100 mg.
Lincomycin base 50 mg.

INDICATIONS

Gastrointestinal and respiratory infections caused by lincomycin and spectinomycin sensitive micro-organisms, like *Campylobacter*, *E. coli*, *Mycoplasma*, *Salmonella*, *Staphylococcus*, *Streptococcus* and *Treponema* spp. in calves, cats, dogs, goats, sheep and swine.

DOSAGE

For intramuscular administration.

Calves:

1 ml per 10 kg body weight for 4 days.

Goats and sheep:

1 ml per 10 kg body weight for 3 days.

Swine:

1 ml per 10 kg body weight for 3 - 7 days.

Cats and dogs:

1 ml per 5 kg body weight for 3 - 5 days, with a maximum of 21 days.

WITHDRAWAL TIMES

- For meat:

Calves, goats, sheep and swine : 14 days.

- For milk

: 3 days.

PACKAGING

Vial of 50 ml and 100 ml.



sales@interchemie.com | interchemie.com

Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XFILSP-EN-01-28082020