

Intertrim LA

Sulfadoxine & Trimethoprim injection



*Gastrointestinal, respiratory and urinary tract infections
caused by trimethoprim and
sulfadoxine sensitive micro-organisms*

Intertrim LA

Solution for parenteral administration

The combination of trimethoprim and sulfadoxine acts synergistic and usually bactericidal against many Gram-positive and Gram-negative bacteria like *E. coli*, *Haemophilus*, *Pasteurella*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. Both compounds affect bacterial purine synthesis in a different way, as a result, of which a double blockade is accomplished.

CONTAINS PER ML

Sulfadoxine 200 mg.
Trimethoprim 40 mg.

INDICATIONS

Gastrointestinal, respiratory and urinary tract infections caused by trimethoprim and sulfadoxine sensitive bacteria, like *E. coli*, *Haemophilus*, *Pasteurella*, *Salmonella*, *Staphylococcus* and *Streptococcus* spp. in calves, cattle, goats, sheep and swine.

DOSAGE

For intramuscular administration.

General:

1 ml per 10 - 15 kg body weight for 3 - 5 days.

WITHDRAWAL TIMES

- For meat : 10 days.
- For milk : 3 days.

PACKAGING

Vial of 100 ml.



sales@interchemie.com | interchemie.com



Manufactured by
Interchemie werken "De Adelaar" B.V.
Metaalweg 8, Venray, The Netherlands

XFISDT-EN-01-28082020