Marbosan-100

Marbofloxacin 10% injection



Treatment of respiratory infections, acute mastitis caused by Gram-negative bacteria in cattle and Metritis Mastitis Agalactia syndrome in pigs.

Marbos

Marbosan-100

Solution for parenteral administration

Marbofloxacin is a synthetic, bactericidal antimicrobial, belonging to the fluoroguinolone group which acts by inhibition of DNA avrase. It has a broad-spectrum activity in vitro against Gram-negative (Pasteurella multocida, Mannheimia haemolytica, Histophilus somni, E. coli) and against Gram-positive bacteria (in particular Staphylococcus). Resistance to Streptococcus may occur.

CONTAINS PER ML

INDICATIONS

Cattle:

Treatment of respiratory infections caused by sensitive strains of Pasteurella multocida, Mannheimia haemolytica, Histophilus somni and Mycoplasma bovis. It is also indicated for the treatment of acute mastitis caused by Gram-negative bacteria susceptible to marbofloxacin.



Pigs:

Treatment of Metritis Mastitis Agalactia syndrome (postpartum dysgalactiae syndrome, PDS) caused by marbofloxacin-sensitive bacterial strain.

DOSAGE

Cattle:

2 mg of marbofloxacin/kg body weight

(corresponding to 1 ml Marbosan-100 per 50 kg body weight) as a subcutaneous or an intramuscular injection once daily for 3 - 5 davs.

The treatment for acute mastitis is 3 days intramuscularly. The first injection may be administered intravenousy.

Pigs:

2 mg of marbofloxacin/kg body weight (corresponding to 1 ml Marbosan-100 per 50 kg body weight) once daily by intramuscular injection for 3 days.

WITHDRAWAL TIMES

Cattle:

- Meat and offal
- : 6 days. - Milk (3-5 days treatment): 36 hours.
- Pigs:
- Meat and offal: 4 days.

PACKAGING

Vial of 100 ml.







sales@interchemie.com | interchemie.com

Manufactured by Interchemie werken "De Adelaar" B.V. Metaalweg 8, Venray, The Netherlands XFIMA10-EN-01-02092020